

Advil®

Relief you can trust

Measure dosing only with the device that comes with the product.



Available in three flavors!
• Fruit • Grape • Blue Raspberry



	Infants' Advil® Concentrated Drops for ages 6-23 months	Children's Advil® Suspension for ages 2-11 years	Advil® Tablets Advil® Liqui-Gels® for ages 12 and up
Active ingredients	Ibuprofen 50 mg (NSAID)* in each 1.25 mL	Ibuprofen 100 mg (NSAID)* in each 5 mL	Ibuprofen 200 mg (NSAID)* in each tablet
Indications	Fever reducer, relieves minor aches and pains due to cold, flu, headaches, and toothaches	Fever reducer, relieves minor aches and pains due to cold, flu, sore throat, headaches, and toothaches	Fever reducer, relieves minor aches and pains due to headache, toothache, backache, menstrual cramps, cold, muscular aches, minor pain of arthritis

		Weight	Age			
		under 6 months		do not use	do not use	do not use
Patient Dosing	12-17 pounds	6-11 months	1.25 mL [†] every 6-8 hrs not to exceed 4 doses/24 hrs	Use Infants' Advil® Concentrated Drops	↓	Use Infants' Advil® Concentrated Drops
	18-23 pounds	12-23 months	1.875 mL [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓	↓
	24-35 pounds	2-3 years	Use Children's Advil® Suspension	1 tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	Use Children's Advil® Suspension
	36-47 pounds	4-5 years	↓	1½ tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
	48-59 pounds	6-8 years	↓	2 tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
	60-71 pounds	9-10 years	↓	2½ tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
	72-95 pounds	11 years	↓	3 tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
		12+ years	Use Advil® Tablets or Liqui-Gels®	Use Advil® Tablets or Liqui-Gels®		1 or 2 tablets [‡] every 4-6 hrs not to exceed 6 tablets/24 hrs

*Nonsteroidal anti-inflammatory drug.

†Measure dosing only with the device provided.

‡If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Please refer to drug facts on the product label for complete warnings.

Important Instructions for Proper Use

- Read and keep the cartons for complete warnings and information on Advil® products
- Remind patients that if they are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), they should consult with you before taking any Advil® products
- Measure dosing only with the device provided
- Adult strength medicines should not be administered to children
- Products contain other ingredients. Please see full ingredient list on each package

Visit advil.com for drug facts, product updates, and additional information.