



## Sigmoidoscopy

Please read these instructions one week prior to your procedure.

### What is a Sigmoidoscopy?

Flexible sigmoidoscopy is a diagnostic procedure used to screen for abnormalities in the sigmoid colon. This procedure examines the sigmoid colon (the lower third of the colon) using a lighted flexible scope that is inserted into the anus and advanced to the splenic flexure. It is possible to take biopsies during this procedure.

### When should I arrive and how long will the procedure take?

Please arrive at the GI Lab 30 minutes prior to the procedure, so that nursing, anesthesia (if applicable) and the physician can evaluate you and prepare you for your procedure. Plan to spend approximately 1 ½-2 hours at the GI Lab.

### Please note:

The time your procedure is scheduled is our best estimation as to when your procedure will begin. Circumstances may arise, ie, emergencies, that may change the scheduled time. We will do our best to notify you as to any changes in the schedule.

### GI Lab Locations

<b>Evanston Hospital</b>	2650 Ridge Ave.	Room 1134	Evanston, IL 60201	(847) 570-2236
<b>Glenbrook Hospital</b>	2100 Pfingsten Road	1 <sup>st</sup> Floor	Glenview, IL 60026	(847) 657-5780
<b>Highland Park Hospital</b>	777 Park Ave West	B345	Highland Park, IL 60035	(847) 480-2828
<b>Highland Park Medical Group Gastroenterology</b>	1777 Green Bay Road	201	Highland Park, IL	(847) 681-4229
<b>Skokie Hospital</b>	9600 Gross Point Road	1 <sup>st</sup> Floor	Skokie, IL 60076	(847) 933-6565
<b>Vernon Hills Specialty Suites</b>	225 N. Milwaukee Ave.	1 <sup>st</sup> Floor	Vernon Hills, IL 60061	(847) 941-7660

### What should I do if I am a diabetic?

Please ask your primary care physician (PCP) to instruct you regarding taking your diabetes medications prior to your Sigmoidoscopy.

### What should I do if I am on blood thinners or have a cardiac stent?

If you have a cardiac stent OR you are on a medication such as Coumadin, Plavix, Ticlid, Lovenox, Pradaxa, Xarelto or any other blood thinner, please contact your prescribing physician to determine

whether you should discontinue this medication prior to the procedure. Notify our office if you are advised not to stop one or more of these medications. It is OK to continue aspirin or other NSAIDs (Advil, Aleve, Ibuprofen, etc) prior to your procedure.

**What should I do if I have a heart defibrillator or pacemaker?**

Your device must be checked within 6 months of your scheduled procedure. If it has not been checked at a NorthShore facility, please bring documentation on the day of your procedure confirming that it has been checked at another facility.

If you have a defibrillator or pacemaker, please contact the GI Lab prior to your scheduled procedure.

Glenbrook	(847) 657-5780
Evanston	(847) 570-2236
Highland Park	(847) 480-2828
Skokie	(847) 933-6565

**❖ Several days prior to the procedure:**

- Purchase two fleets enemas in any drug store or pharmacy without a prescription.
- Check with your insurance company to see if you need to be pre-certified before the procedure or if you need to get a referral from your primary care physician.

**❖ On the Day of Your Procedure:**

- Take 1st fleets enema 2 hours before you are to arrive in the GI Lab.
- Take 2nd fleets enema 1 hour before you are to arrive in the GI Lab. You should lie on your left side when giving yourself the enema. To get best results, try to hold it in for as long as possible (i.e. 5 minutes)
- If your procedure is scheduled in the morning, only have clear liquids for breakfast.
- If your procedure is scheduled in the afternoon, have a light breakfast and clear liquids for lunch.